

Climate Drift in CCSM4 Decadal Prediction Experiments

Haiyan Teng
NCAR

Toulouse, May 14, 2013

**Grant Branstator, Jerry Meehl, Alicia Karspeck, Steve Yeager,
Gokhan Danabasglu, Joe Tribbia, DART, CSEG ...**

CCSM4 Decadal Prediction Experiments

		Hindcast	DART
IC	atm/lnd	20thC, rcp4.5	AMIP
	ocn	Hindcast	DART
	ice	Hindcast (4-yr salinity restoring)	Hindcast (3-mon salinity restoring)
start years		61,66,71,76,81,86,91,96 00,01,02,03,04,05,06 (15 total)	75,80,85,90,95 00,01,02,03,04,05,06 (12 total)
ensemble#		10	
length		10/30 years	


Free runs: 20thC+RCP4.5, 6 members

last 700 yrs of the 1300-yr preindustrial control


Definition of drift:

Initialized decadal prediction minus the **uninitialized** (20thC +RCP4.5)

Drift from a particular start year is caused by model climatological bias and initial anomalies



Motivation











How long does the model drift?
=> how long impact from initialization can last


What cause the drift?

SST


Precipitation



Drift in global avg TAS and AMOC







SST: decadal prediction minus free



Nino34


Global TAS



- 1961
 - 1966
 - 1971
 - 1976/1975
 - 1981/1980
 - 1986/1985
 - 1991/1990
 - 1996/1995
 - 2000
 - 2001
 - 2002
 - 2003
 - 2004
 - 2005
 - 2006
 - avg
- HdI
- HdII


Drift in DART DJF SLP

Highly connected to drift in eq. Pacific SST




AMOC is still drifting


Eq. Pacific TEMP Diffs in month 1








What caused the drift?




Nino34 region (60m) heat budget






Unusual initial conditions

can cause the extended cold drift







Zonal current (2.5S-2.5N)



Nino34

Global TAS




- 1961
 - 1966
 - 1971
 - 1976/1975
 - 1981/1980
 - 1986/1985
 - 1991/1990
 - 1996/1995
 - 2000
 - 2001
 - 2002
 - 2003
 - 2004
 - 2005
 - 2006
 - avg
- HdI
- HdII


Summary

- The Hindcast initial conditions cause a warm drift from 1961, 1966, 1971, and 1976 start years, and the other Hindcast initial conditions as well as those from DART cause a cold drift in the equatorial Pacific SST in the first five prediction years.
- The warm drift is mainly caused by large initial warm anomalies at thermocline and associated feedbacks, while the strong upwelling and zonal current anomalies are two major players for the extended equatorial cold drift.
- The warm/cold drifts may contaminate the prediction anomalies (drift $\neq$ monotonic decay), adding difficulty to interpret the prediction result.


Implications for decadal prediction




Influence from initial information will last longer





Hindcast I





Hindcast II





DART

 $^{\circ}\text{C}$

Nino34




Global TAS



But why DART 2003-2006 initial conditions also produce colder anomalies than Uninit?

SST score (correlation)



Stippling represents 95% sig lev where Hindcast/DART score exceeds AR1 model (built with lag1 from the 700-yr control, trend from the uninit, and noise)

- Score is low in the eq. Pacific due to dominance of ENSO (noise).
- The low score in the eq. Pacific does not exclude possibility that small components of the eq. Pacific SST (e.g. associated with IPO) may be more predictable...(same Idea as Branstator's AMOC talk).